

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

878-082  
09/02/98

# WEST

## Freeform Search

Database:

US Patents Full-Text Database  
 US Pre-Grant Publication Full-Text Database  
 JPO Abstracts Database  
 EPO Abstracts Database  
 Derwent World Patents Index  
 IBM Technical Disclosure Bulletins

Term:

L4 and fluorescence

Display:  Documents in Display Format:  Starting with Number Generate: ☐ Hit List ☒ Hit Count ☐ Side by Side ☐ Image

Search

Clear

Help

Logout

Interrupt

Main Menu

Show S Numbers

Edit S Numbers

Preferences

Cases

---

### Search History

---

DATE: Monday, June 24, 2002   [Printable Copy](#)   [Create Case](#)

Set Name   Query  
 side by side

Hit Count   Set Name  
                     result set

DB=USPT,JPAB,EPAB,DWPI; PLUR=YES; OP=ADJ

<u>L5</u>	L4 and fluorescence	9	<u>L5</u>
<u>L4</u>	L3 and alle\$ specific primer\$1	13	<u>L4</u>
<u>L3</u>	L2 and sequencing	441	<u>L3</u>
<u>L2</u>	monitor\$ near5 amplif\$	10643	<u>L2</u>
<u>L1</u>	monitor\$ nea5 amplif\$ near5 sequencing	0	<u>L1</u>

END OF SEARCH HISTORY

Generate Collection

Print

**Search Results - Record(s) 1 through 9 of 9 returned.**

- 
- ☐ 1. 6391558. 14 Apr 00; 21 May 02. Electrochemical detection of nucleic acid sequences. Henkens; Robert W., et al. 435/6; 422/50 422/62 422/63 422/67 422/68.1 422/69 422/82.01 435/91.1 435/91.2. C12Q001/68 C12P019/34 G01N015/06 G01N030/96 G01N027/00.
- 
- ☒ 2. 6361949. 22 Feb 00; 26 Mar 02. Nucleic acid amplification with direct sequencing. Sommer; Steven Seev. 435/6; C12Q001/68 C12Q001/70.
- 
- ☒ 3. 6322976. 17 Mar 99; 27 Nov 01. Compositions and methods of disease diagnosis and therapy. Aitman; Timothy J., et al. 435/6; 435/7.23 536/23.1 536/24.3 536/24.31. C12Q001/68 G01N033/574 C07H021/02 C07H021/04 C07H021/00.
- 
- ☒ 4. 6316198. 18 Mar 00; 13 Nov 01. Detection of mutations in genes by specific LNA primers. Skouv; Jan, et al. 435/6; 435/91.1 435/91.2 536/23.1 536/24.3. C12Q001/68 C12P019/34 C07H021/02.
- 
- ☐ 5. 6117635. 11 Apr 97; 12 Sep 00. Nucleic acid amplification oligonucleotides with molecular energy transfer labels and methods based thereon. Nazarenko; Irina A., et al. 435/6; 435/91.2 536/22.1 536/24.33 536/25.32. C12Q001/68 C12P019/34 C07H021/04 C07H021/00.
- 
- ☒ 6. 6090552. 11 Jul 97; 18 Jul 00. Nucleic acid amplification oligonucleotides with molecular energy transfer labels and methods based thereon. Nazarenko; Irina A., et al. 435/6; 435/91.2 536/24.3 536/24.32 536/24.33. C12Q001/68 C12P019/34 C07H021/04 C12N015/00.
- 
- ☐ 7. 6027913. 27 Dec 94; 22 Feb 00. Nucleic acid amplification with direct sequencing. Sommer; Steven S.. 435/69.1; 435/91.21. C12P021/00 C12P019/34.
- 
- ☒ 8. 6015670. 14 Nov 97; 18 Jan 00. Methods for identifying a mutation in a gene of interest without a phenotypic guide using ES cells. Goodfellow; Peter N.. 435/6; 435/91.2. C12Q001/68 C12P019/34.
- 
- ☐ 9. 5994075. 16 May 97; 30 Nov 99. Methods for identifying a mutation in a gene of interest without a phenotypic guide. Goodfellow; Peter N.. 435/6; 435/441 435/444 435/446 435/91.2. C12Q001/68 C12P019/34.
- 

Generate Collection

Print

Term	Documents
FLUORESCENCE.DWPI,EPAB,JPAB,USPT.	53629
FLUORESCENCES.DWPI,EPAB,JPAB,USPT.	358
(4 AND FLUORESCENCE).USPT,JPAB,EPAB,DWPI.	9
(L4 AND FLUORESCENCE).USPT,JPAB,EPAB,DWPI.	9

[Previous Page](#)[Next Page](#)

STN much 09/818086

=> s monitor?(10a)amplif?(10a)fluorescen?  
L1 92 MONITOR?(10A) AMPLIF?(10A) FLUORESCEN?

=> s l1 and sequencing  
L2 5 L1 AND SEQUENCING

=> s l2 and allele? specific primer#  
L3 0 L2 AND ALLEL? SPECIFIC PRIMER#

=> s l1 and allele? specific primer#  
L4 2 L1 AND ALLEL? SPECIFIC PRIMER#

=> s l4 and sequencing  
L5 0 L4 AND SEQUENCING

=> d l4 1-2 bib ab

L4 ANSWER 1 OF 2 MEDLINE  
AN 2000115340 MEDLINE  
DN 20115340 PubMed ID: 10649496  
TI Mutation detection by TaqMan-allele specific amplification: application to molecular diagnosis of glycogen storage disease type Ia and medium-chain acyl-CoA dehydrogenase deficiency.  
AU Fujii K; Matsubara Y; Akanuma J; Takahashi K; Kure S; Suzuki Y; Imaizumi M; Iinuma K; Sakatsume O; Rinaldo P; Narisawa K  
CS Department of Medical Genetics, Tohoku University School of Medicine, Sendai, Japan.  
SO HUMAN MUTATION, (2000) 15 (2) 189-96.  
Journal code: 9215429. ISSN: 1059-7794.  
CY United States  
DT Journal; Article; (JOURNAL ARTICLE)  
LA English  
FS Priority Journals  
EM 200002  
ED Entered STN: 20000309  
Last Updated on STN: 20000309  
Entered Medline: 20000224  
AB We have devised an allele-specific amplification method with a TaqMan fluorogenic probe (TaqMan-ASA) for the detection of point mutations. Pairwise PCR amplification using two sets of **allele-specific primers** in the presence of a TaqMan probe was **monitored** in real time with a **fluorescence** detector. Difference in **amplification** efficiency between the two PCR reactions was determined by "threshold" cycles to differentiate mutant and normal alleles without post-PCR processing. The method measured the efficiency of amplification rather than the presence or absence of end-point PCR products, therefore allowing greater flexibility in designing **allele-specific primers** and an ample technical margin for allelic discrimination. We applied the TaqMan-ASA method to detect a prevalent 727G>T mutation in Japanese patients with glycogen storage disease type Ia and a common 985A>G mutation in Caucasian patients with medium-chain acyl-CoA dehydrogenase deficiency. The method can be automated and may be applicable to the DNA diagnosis of various genetic diseases.  
Copyright 2000 Wiley-Liss, Inc.

L4 ANSWER 2 OF 2 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.  
AN 2000:113844 BIOSIS  
DN PREV200000113844  
TI Mutation detection by TaqMan-allele specific amplification: Application to molecular diagnosis of glycogen storage disease type Ia and medium-chain acyl-CoA dehydrogenase deficiency.  
AU Fujii, Kunihiro; Matsubara, Yoichi (1); Akanuma, Jun; Takahashi, Kazutoshi; Kure, Shigeo; Suzuki, Yoichi; Imaizumi, Masue; Iinuma, Kazuie;

Sakatsume, Osamu; Rinaldo, Piero; Narisawa, Kuniaki  
 CS (1) Department of Medical Genetics, Tohoku University School of Medicine,  
 1-1 Seiryomachi, Aobaku, Sendai, 980-8574 Japan  
 SO Human Mutation, (2000) Vol. 15, No. 2, pp. 189-196.  
 ISSN: 1059-7794.  
 DT Article  
 LA English  
 SL English  
 AB We have devised an allele-specific amplification method with a TaqMan  
 fluorogenic probe (TaqMan-ASA) for the detection of point mutations.  
 Pairwise PCR amplification using two sets of **allele-**  
**specific primers** in the presence of a TaqMan probe was  
**monitored** in real time with a **fluorescence** detector.  
 Difference in **amplification** efficiency between the two PCR  
 reactions was determined by "threshold" cycles to differentiate mutant and  
 normal alleles without post-PCR processing. The method measured the  
 efficiency of amplification rather than the presence or absence of  
 end-point PCR products, therefore allowing greater flexibility in  
 designing **allele-specific primers** and an  
 ample technical margin for allelic discrimination. We applied the  
 TaqMan-ASA method to detect a prevalent 727G>T mutation in Japanese  
 patients with glycogen storage disease type Ia and a common 985A>G  
 mutation in Caucasian patients with medium-chain acyl-CoA dehydrogenase  
 deficiency. The method can be automated and may be applicable to the DNA  
 diagnosis of various genetic disease.

=> dup rem l2  
 PROCESSING COMPLETED FOR L2  
 L6 5 DUP REM L2 (0 DUPLICATES REMOVED)

=> d l6 1-5 bib ab kwic

L6 ANSWER 1 OF 5 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.  
 AN 2001:334723 BIOSIS  
 DN PREV200100334723  
 TI Methods and devices for hemogeneous nucleic acid amplification and  
 detector..  
 AU Higuchi, Russell G. (1)  
 CS (1) San Francisco, CA USA  
 ASSIGNEE: Roche Molecular Systems, Inc.  
 PI US 6171785 January 09, 2001  
 SO Official Gazette of the United States Patent and Trademark Office Patents,  
 (Jan. 9, 2001) Vol. 1242, No. 2, pp. No Pagination. e-file.  
 ISSN: 0098-1133.  
 DT Patent  
 LA English  
 AB This invention relates to improved methods for nucleic acid detection  
 using methods such as the polymerase chain reaction (PCR). More  
 specifically, the invention provides methods for simultaneous  
 amplification and detection to enhance the speed and accuracy of prior  
 methods. The methods involve the introduction of detectable DNA binding  
 agents into the amplification reaction, which agents produce a detectable  
 signal that is enhanced upon binding double-stranded DNA. In a preferred  
 embodiment, the binding agent is a **fluorescent** dye. The methods  
 also provide means for **monitoring** the increase in product DNA  
 during an **amplification** reaction.  
 AB. . . produce a detectable signal that is enhanced upon binding  
 double-stranded DNA. In a preferred embodiment, the binding agent is a  
**fluorescent** dye. The methods also provide means for  
**monitoring** the increase in product DNA during an  
**amplification** reaction.  
 IT Methods & Equipment  
 polymerase chain reaction [PCR]: DNA amplification, DNA amplification

method, detection method, in-situ recombinant gene expression  
detection, **sequencing** techniques, simultaneous  
amplification-detection

L6 ANSWER 2 OF 5 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

AN 2001:317029 BIOSIS

DN PREV200100317029

TI **Monitoring** of mixed chimerism by a technique using  
**fluorescence** based PCR **amplification** of microsatellite  
after allogeneic hematopoietic stem cell transplantation.

AU Saito, Akiko (1); Ogawa, Seishi (1); Hadama, Tohru; Kinoshita, Moritoshi;  
Chiba, Shigeru (1); Hirai, Hisamaru (1)

CS (1) Hematology and Oncology, University of Tokyo, Bunkyo-ku, Tokyo Japan

SO Blood, (November 16, 2000) Vol. 96, No. 11 Part 1, pp. 395a. print.

Meeting Info.: 42nd Annual Meeting of the American Society of Hematology  
San Francisco, California, USA December 01-05, 2000 American Society of  
Hematology

. ISSN: 0006-4971.

DT Conference

LA English

SL English

AB (Introduction) Monitoring of mixed chimerism following hematopoietic stem  
cell transplantation (HSCT) provides an important clue to evaluate  
engraftment and to detect graft failure or early relapse. Several  
techniques have been applied for this purpose; Mixed chimerism after  
sex-mismatched transplant can be quickly and quantitatively assessed by  
fluorescent in situ hybridization (FISH) analysis using X- and Y-specific  
probes. Assessment of chimerism in sex-matched transplant has also been  
possible by differentially detecting a polymorphic allele(s) between the  
donor and recipient. However, the conventional methods for quantitative  
detection of polymorphisms such as VNTRs have been frequently too  
time-consuming in the context of clinical applications. In this study we  
intended to develop a simple method for quickly estimating post-transplant  
chimerism. (Materials and methods) Genomic DNA was extracted from bone  
marrow and/or blood samples of 27 donor-recipient pairs following  
allogeneic HSCT and subjected to the microsatellite PCR analysis, in which  
three microsatellite loci, D18S51, D20S471 and D22S684, were PCR-amplified  
using fluorescent primers from the genomic DNAs and length of the PCR  
products were analyzed using an ABI PRISM 377 automated sequence analyzer.  
Because the polymorphism in a given locus is represented by the difference  
in the length of the corresponding PCR products, we first determined the  
informative loci which showed different electrophoretic mobilities between  
the donor-recipient pair, and then assessed the chimerism in a given  
sample by measuring relative intensity of each polymorphic peak for the  
informative loci. Reliability of this assay was tested by measuring  
chimerism of the standard DNA samples whose donor/recipient-composition  
was already known, and by comparing the results with those obtained from  
other assays, for example, XY-FISH. (Results) In our method, 11 of 11  
(100%) cases transplanted from unrelated donors and 13 of 16 (81%) cases  
from related donors had at least one informative microsatellite locus.  
Measurement of the standard DNA samples show a linear correlation between  
the measured values for donor-recipient ratios and the standardized values  
for the DNA composition. More than 10% of chimera can be stably detected,  
using as little as ten nanograms of sample DNA. In 11 patients, results  
from the microsatellite PCR showed excellent concordance with the data  
obtained from the conventional FISH analysis using X- and Y-specific  
probes and/or probes detecting tumor-specific translocations.  
(Conclusions) Fluorescent primer-based microsatellite PCR assay is a  
feasible, rapid and reliable technique for assessment of mixed chimerism  
after allogeneic HSCT, even with minuscule samples.

TI **Monitoring** of mixed chimerism by a technique using  
**fluorescence** based PCR **amplification** of microsatellite  
after allogeneic hematopoietic stem cell transplantation.

IT . . .

hybridization [FISH]: diagnostic method; microsatellite PCR  
[microsatellite polymerase chain reaction]: DNA amplification,  
amplification method, fluorescence-based, in-situ recombinant gene  
expression detection, **sequencing** techniques

IT Miscellaneous Descriptors

chromosomal translocations: tumor-specific; electrophoretic mobility;  
engraftment; mixed chimerism; Meeting Abstract; Meeting Poster

L6 ANSWER 3 OF 5 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

AN 1999:135568 BIOSIS

DN PREV199900135568

TI Identification of *Leptospira biflexa* by real-time homogeneous detection of  
rapid cycle PCR product.

AU Woo, T. H. S; Patel, B. K. C. (1); Cinco, M.; Smythe, L. D.; Norris, M.  
A.; Symonds, M. L.; Dohnt, M. F.; Piispanen, J.

CS (1) Sch. Biomol. Biomed. Sci., Fac. Sci., Griffith Univ., Nathan Campus,  
Brisbane, QLD 4111 Australia

SO Journal of Microbiological Methods, (Feb., 1999) Vol. 35, No. 1, pp.  
23-30.

ISSN: 0167-7012.

DT Article

LA English

AB Sequence analysis of 16S rRNA genes extracted from nucleic acids databases  
enabled the identification of a *Leptospira biflexa* (*L. biflexa*) signature  
sequence, against which a reverse primer designated L613, was designed.  
This primer, when used in conjunction with a universal bacterial specific  
forward primer designated Fd1, enabled the development of a  
LightCycler<sup>TM</sup>-based PCR protocol in which **fluorescence** emission  
due to binding of SYBR Green I dye to **amplified** products could  
be detected and **monitored**. A melting temperature ( $T_m$ ),  
determined from the melting curve of the amplified product immediately  
following the termination of thermal cycling, confirmed that the product  
was that of *L. biflexa*. Agarose gel electrophoresis therefore was not  
necessary for identification of PCR products. The PCR protocol was very  
rapid, and consisted of 30 cycles with a duration of 20 s for each cycle  
with the monitoring of the melting curve requiring an additional 3 min.  
The whole protocol was completed in less than 20 min. The PCR protocol was  
also specific and enabled the identification of 18 strains of *L. biflexa*,  
whilst excluding 14 strains of *L. interrogans* and *Leptonema illini*. Two  
examples of its utility in improving work flow of a *Leptospira* reference  
laboratory are presented in this article. The use of a simple boiling  
method for extraction of DNA from all the members of the Leptospiraceae  
family DNA further simplifies the procedure and makes its use conducive to  
diagnostic laboratories.

AB. . . conjunction with a universal bacterial specific forward primer  
designated Fd1, enabled the development of a LightCycler<sup>TM</sup>-based PCR  
protocol in which **fluorescence** emission due to binding of SYBR  
Green I dye to **amplified** products could be detected and  
**monitored**. A melting temperature ( $T_m$ ), determined from the melting  
curve of the amplified product immediately following the termination of  
thermal cycling, . . .

IT

. . . analytical method, gel electrophoresis; DNA extraction:  
Isolation/Purification Techniques: CB, extraction method; LightCycler  
PCR [polymerase chain reaction]: DNA amplification, amplification  
method, **sequencing** techniques, in-situ recombinant gene  
expression detection

IT Miscellaneous Descriptors

nucleotide sequence

L6 ANSWER 4 OF 5 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.

AN 1999:219054 BIOSIS

DN PREV199900219054

TI Continuous fluorescent monitoring of rapid cycle polymerase chain

reaction.

- AU Pritham, Gregory H.; Wittwer, Carl T. (1)  
CS (1) Department of Pathology, University of Utah Medical School, 50 N.  
Medical Drive, Salt Lake City, UT, 84132 USA  
SO Journal of Clinical Ligand Assay, (Winter, 1998) Vol. 21, No. 4, pp.  
404-412.  
ISSN: 1081-1672.  
DT Article  
LA English  
SL English  
AB Polymerase chain reaction (PCR) amplification and analysis can be  
performed rapidly. Indeed, both amplification and analysis can occur  
simultaneously in the same instrument in only 10-30 minutes. Rapid cycle  
PCR is possible because denaturation, annealing, and extension are fast  
reactions. Currently, cycling speeds are limited by instrumentation, not  
chemistry. If rapid cycle PCR is continuously monitored with a  
fluorimeter, amplification progress can be followed with double-stranded  
DNA specific dyes or resonance energy transfer probes of multiple designs.  
Initial template copy number can be determined by monitoring fluorescence  
once each cycle. Continuous monitoring of fluorescence within a cycle as  
the temperature is changing can be used to follow product or probe  
hybridization. Fluorescence melting curves immediately after amplification  
provide dynamic dot blots of hybridization for product identification or  
single base genotyping.
- IT Methods & Equipment  
fluorimeter: laboratory equipment; genotyping: analytical method;  
polymerase chain reaction: DNA amplification, analytical method,  
**sequencing** techniques, molecular genetic method, in-situ  
recombinant gene expression detection; rapid cycle polymerase chain  
reaction-continuous **fluorescent monitoring**: DNA  
**amplification, sequencing** techniques, molecular  
genetic method, analytical method, in-situ recombinant gene expression  
detection
- IT Miscellaneous Descriptors  
instrumentation; melting curves; mutations: detection; template. . .
- L6 ANSWER 5 OF 5 BIOSIS COPYRIGHT 2002 BIOLOGICAL ABSTRACTS INC.  
AN 1998:472877 BIOSIS  
DN PREV199800472877  
TI Fluorescence cross-correlation: A new concept for polymerase chain  
reaction.
- AU Rigler, Rudolf (1); Foeldes-Papp, Zeno; Meyer-Almes, Franz-Josef; Sammet,  
Cyra; Voelcker, Martin; Schnetz, Andreas  
CS (1) Dep. Med. Biophys., MBB, Karolinska Inst., S-17177 Stockholm Sweden  
SO ~~Journal of Biotechnology~~, (Aug. 12, 1998) Vol. 63, No. 2, pp. 97-109.  
ISSN: 0168-1656.  
DT Article  
LA English  
AB In this article we present a new concept for the detection of any  
specifically amplified target DNA sequences in multiple polymerase chain  
reactions (PCR) based on fluorescence correlation spectroscopy (FCS). The  
accumulation of double-stranded target DNA is **monitored** by the  
cross-correlated **fluorescence** signals provided by two  
**amplification** primers which are 5'-tagged with two different kinds  
of fluorophores (Rhodamine-Green and Cy5). Only the amplified target DNA  
sequence carrying both primers is observed. Its signal emerges from the  
background of non-incorporated or non-specifically incorporated primers.  
Down to 10-25 initial copy numbers of the template in the PCR compartment  
DNA can presently be detected. No external or internal standards are  
required for determining the size and the amplified copy number of  
specific DNA. The PCR amplification process is started with all  
ingredients in a single compartment (e.g. of a microtiter plate), in which  
amplification and measurement are performed. This eliminates the need for  
post-PCR purification steps. The homogeneous one-tube approach does not



depend on fluorescence energy transfer between the fluorogenic dyes. Thus, it does not interfere with the enzymatic amplification reaction of PCR and allows the continued use of different conditions for amplifying DNA. The results exemplified by PCR-amplified 217-bp and 389-bp target DNA sequences demonstrate that the analysis based on two-color fluorescence cross-correlation is a powerful method for simplifying the identification of targets in PCR for medical use. For this purpose, an instrument optimized for two-color excitation and detection of two-color emission has been developed, incorporating the principle of confocal arrangement.

AB. . . sequences in multiple polymerase chain reactions (PCR) based on fluorescence correlation spectroscopy (FCS). The accumulation of double-stranded target DNA is **monitored** by the cross-correlated **fluorescence** signals provided by two **amplification** primers which are 5'-tagged with two different kinds of fluorophores (Rhodamine-Green and Cy5). Only the amplified target DNA sequence carrying. . .

IT Methods & Equipment

polymerase chain reaction: DNA amplification, in-situ recombinant gene expression detection, **sequencing** techniques, molecular genetic method; two-color fluorescence cross-correlation spectroscopy: analytical method

IT Miscellaneous Descriptors

biotechnology

=>